

TECHNICAL DATASHEET


IOCID

Zootechny / Teat hygiene

Post-dipping gel with povidone iodine

Features

Post-milking gel with povidone iodine (1150 ppm). Thanks to its formulation, IOCID helps protect the breasts and performs an emollient and soothing action while preserving the natural characteristics of the skin. IOCID has a stable color which always allows a clear identification of the treated parts.



Application fields

IOCID is employed in the zootechny-veterinary sector.

How to use

Treat the affected parts directly using the appropriate applicator cup. Before milking, clean the treated parts thoroughly with paper or disposable wipes.

Concentration of use: as it is, without dilution

Leave to act for: until the next milking

Features

Form:	gel
pH:	4-5
density:	1,01 – 1,05 g/cm ³ @25 °C

Compatibility

No incompatibility known.

Packaging

Canister: 10 kg 20 kg

Barrel: 200 kg

IBC: 1000 kg

Other information

IOCID does not meet the classification criteria of Regulation (EC) no. 1272/2008 (CLP).

IOCID is not defined as dangerous for the different modes of transport according to the "United Nations Recommendations on the Transport of Dangerous Goods - Model Regulations".



AZIENDA CERTIFICATA
SECONDO GLI STANDARD
DI QUALITÀ ISO 9001:2015

Attention

This technical data sheet is intended to be purely indicative. For the definition of the correct protocol for the use of Devidet s.r.l. always consult the Devidet s.r.l. Technical Office, expert technicians will be able to provide the most suitable solutions for specific needs. Product for professional use only. We decline all responsibility for damage resulting from improper or unauthorized use of the product. Before use, check that the product matches the type of application required. Before use, consult the safety data sheet or the information sheet.

Revision 10 16/01/2021

